

# QA0-FR6-1000

## FragIT™ MaxiSpin Columns Instructions for Use

Cleave 10-100mg IgG per column

---

### Product Description

FragIT™ MaxiSpin column contains FabRICATOR® enzyme covalently coupled to agarose beads for cleavage of IgG to pure F(ab')<sub>2</sub> and Fc fragments. IgG is incubated with the FabRICATOR® agarose beads for 20 minutes. F(ab')<sub>2</sub> and Fc fragments are then collected in a centrifugation step. Since FabRICATOR® enzyme is immobilized on the agarose beads, there is no need for further purification to remove the enzyme from the cleavage products.

FabRICATOR® is a digestive enzyme that cleaves IgG at a specific site below the hinge region yielding pure F(ab')<sub>2</sub> and Fc fragments. Since FabRICATOR® cleaves only at one specific site, there is no risk of getting fragments other than F(ab')<sub>2</sub> and Fc even if the incubation time is increased.

FabRICATOR® cleaves all subclasses of human, monkey, rabbit, and sheep IgG but only mouse IgG2a and IgG3 subclasses. Cleavage of mouse IgG2a requires significantly longer incubation time as compared to cleavage of human IgG and may need to be optimized.

### Contents and Storage

FragIT™ MaxiSpin column is supplied in 20% EtOH. No preservatives are added. One FragIT™ MaxiSpin column is sufficient to cleave 10-100mg IgG.

FragIT™ MaxiSpin is shipped on cold packs and should be stored at 4-8°C upon arrival.

FragIT™ Spin Columns are for *in vitro* R&D use only.

### Quality Control

FabRICATOR® enzyme is tested to ensure lot-to-lot consistency.

FabRICATOR® enzyme is tested for sterility on blood agar plates, Sabaraud dextrose agar plates, and in thioglycolate broth.

FragIT™ Spin Columns are tested to ensure lot-to-lot consistency.

### Additional Materials Required / Not Provided

- ✓ Cleavage buffer: 50mM sodium phosphate, 150mM NaCl, pH 6.6
- ✓ Collection tubes: 50ml centrifuge tubes

### Procedure – IgG Cleavage

★ **Make sure your antibody is in cleavage buffer.**

1. Break off the bottom plastic cap of the spin column and slightly open the screw cap lid 90° counter-clockwise. **SAVE THE BOTTOM CAP!** Lids and bottom caps are used during the incubation step.
2. Place the spin column in a 50ml centrifuge tube.
3. Centrifuge the spin column at 100 x g for one (1) minute to remove storage solution.
4. Equilibrate the spin column with 10.0ml of cleavage buffer.
5. Centrifuge the spin column at 100 x g for one (1) minute.
6. Repeat steps 4 and 5 two more times.
7. Re-insert the bottom cap on the spin column and seal tightly.
8. Immediately add 5.0-10.0ml of IgG at a maximum concentration of 20mg/ml in cleavage buffer.
9. Re-seal the spin column with the lid.
10. Take care to stir and suspend the agarose beads manually and make sure the spin column will flow.

11. Incubate the spin column by end-over-end mixing at room temperature for 45 minutes. The incubation time can be increased without over digestion of the IgG. For cleavage of mouse IgG2a, the incubation time may need to be increased to 6 hours.
12. Open the spin column lid and remove the bottom cap.
13. Place the spin column in a 50ml centrifuge tube.
14. Centrifuge the spin column at 100 x g for one (1) minute to elute the sample.

***For maximum recovery of sample:***

15. Add an additional 5ml of cleavage buffer to the spin column.
16. Place the spin column in a 50ml centrifuge tube.
17. Centrifuge the spin column at 100 x g for one (1) minute to elute the sample.
18. Repeat steps 15-17 one more time.

***Product Reference***

Ryan, MH et al. 2008 Proteolysis of purified IgGs by human and bacterial enzymes in vitro and the detection of specific proteolytic fragments of endogenous IgG in rheumatoid synovial fluid. *Molecular Immunology* 45: 1837-1846.