

# **Anti-Hsp70 Monoclonal Antibody**

## ORDERING INFORMATION

**Catalog No.:** 11118 (clone 5A5) **Size:** 100ug Protein G-purified antibody in PBS, pH 7.4 and 50% glycerol.

# **BACKGROUND**

Hsp70 binds ATP with high affinity and possesses a weak ATPase activity which can be stimulated by binding to unfolded proteins and synthetic peptides. Hsp70 recognizes and binds to nascent polypeptide chains as well as partially folded intermediates of proteins preventing their aggregation and misfolding. The binding of ATP triggers a critical conformational change leading to the release of the bound protein. The ability of Hsp70 to undergo cycles of binding and release from hydrophobic stretches of partially unfolded proteins is the basis for their role in a variety of intracellular functions such as protein synthesis, protein folding and oligomerization and protein transport.

# **SPECIFICATION SUMMARY**

Antigen: Recombinant human Hsp70

expressed in *E. coli*. **Host Species:** Mouse

Subclass: IgG1

Preservatives: 0.09% sodium azide

#### **SPECIFICITY**

This antibody recognizes human, mouse, rat, fish, avian, rabbit, amphibian, *Drosophila*, and yeast Hsp70, Hsc70, Grp78, and, following heat shock, Hsp72.

# <u>APPLICATIONS</u>

Immunoblotting: use at 1ug/ml (ECL). A band of ~70 kDa is detected.
Immunoprecipitation: use at 1ug/ml
Immunohistochemistry/Immunofluorescence: use at 2 ug/ml
User should determine optimal
concentrations for their applications.
Positive control: Heat-shocked HeLa
cell lysate.

# **DILUTION INSTRUCTIONS**

Dilute in PBS or medium which is identical to that used in the assay system.

## STORAGE AND STABILITY

This antibody is stable for at least one (1) year at -20°C.

For in vitro investigational use only. Not for use in therapeutic or diagnostic procedures.