

ECS (DDDDK) Affinity-Purified Polyclonal Antibody, Agarose Immobilized

ORDERING INFORMATION

Catalog No.	Size
18830A	100ug

Format: 0.1mg antibody/0.4ml of 50% gel slurry in Phosphate Buffered Saline (PBS), 0.09% sodium azide. Antibody coupled to agarose beads with cyanogen bromide.

BACKGROUND

Enterokinase, also known as enteropeptidase, is a serine protease that catalyzes the hydrolysis of peptide bonds in proteins but, unlike other kinases, does not catalyze transfer of phosphate groups. Enterokinase cleaves proteins following a lysine at an enterokinase cleavage site, or ECS (Asp-Asp-Asp-Asp-Lys). *In vivo*, this cleavage results in trypsin-dependent activation of other pancreatic zymogens, such as chymotrypsinogen, proelastase, procarboxypeptidase and prolipase in the lumen of the gut.

SPECIFICATION SUMMARY

Antigen: DDDDK conjugated to KLH.

Host Species: Goat

Specificity: Antibody was affinity-purified on the peptide.

APPLICATIONS

Immunoprecipitation: use 15-25ul of gel slurry per 0.1-1mg of protein extract or lysate.

These are recommended concentrations. Endusers should determine optimal concentrations for their applications.

DILUTION INSTRUCTIONS

Dilute in PBS or medium that is identical to that used in the assay system.

STORAGE AND STABILITY

This antibody is stable for at least one (1) year at 2-8°C.

For in vitro investigational use only. Not for use in therapeutic or diagnostic procedures.