

TARPγ2/Stargazin Monoclonal Antibody

ORDERING INFORMATION

Catalog No.: 11571 (clone S245-1) Size: 100ug in PBS, pH 7.4; 50% glycerol, 0.09% sodium azide. Purified by Protein G affinity chromatography.

BACKGROUND

Stargazin belongs to a group of voltagegated calcium channels that are present in most excitable cells. It appears to stabilize calcium channels in an inactivated (closed) state. Voltage-gated calcium channels exert spatial and temporal control over cellular calcium concentrations and serve to modulate neurotransmitter release, hormone secretion, muscle contraction, electrical activity, cell metabolism and proliferation, gene expression, and neuronal survival.

SPECIFICATION SUMMARY

Antigen: Fusion protein corresponding to aa 203-323 (cytoplasmic C-terminus) of rat TARPγ2 (accession no. Q71RJ2).

Host Species: Mouse **Antibody Class**: IgG2a

SPECIFICITY

This antibody recognizes mouse and rat TARPγ2. It does not cross-react with TARPγ3, TARPγ4, or TARPγ8.

APPLICATIONS

<u>Immunoblotting</u>: use at 1ug/ml. Predicted molecular weight is 36kDa.

75— 50— 37—

<u>Positive control</u>: Rat brain lysate. These are recommended concentrations. User should determine optimal concentrations for their application.

DILUTION INSTRUCTIONS

Dilute in PBS or medium which is identical to that used in the assay system.

STORAGE AND STABILITY

This antibody is stable for at least one (1) year at -20°C. Avoid repeated freezing and thawing.

For in vitro investigational use only. Not for use in therapeutic or diagnostic procedures.