

# c-myc Affinity-Purified Polyclonal Antibody, HRP conjugate

#### ORDERING INFORMATION

Catalog No. Size 18826P 100ug

**Format:** 1mg/ml conjugated to horseradish peroxidase (HRP) in Phosphate Buffered Saline, pH 7.2, 0.2% BSA, 0.1% Pro-Clean 400. Molar enzyme/antibody ratio = 4:1.

### **BACKGROUND**

c-myc is an epitope tag often constructed on the N- or C-terminus of a protein to allow for detection and analysis of the protein. The peptide sequence of the myc tag is EQKLISEEDL (Glu-Gln-Lys-Leu-Ile-Ser-Glu-Glu-Asp-Leu). The c-myc tag is approximately 1202 Daltons in atomic mass and includes 10 amino acids.

### **SPECIFICATION SUMMARY**

Antigen: Amino acids 410-419 (EQKLISEEDL) of human myc conjugated to KLH.

Host Species: Rabbit

**Specificity:** Antibody was affinity-purified on the peptide.

#### **APPLICATIONS**

*Immunoblotting:* use at a dilution of 1:1,000-1:30,000.

Immunocytochemistry: use at a dilution of

1:200-1:500.


ELISA: use at a dilution of 1:10,000-

1:100,000.

These are recommended dilutions.

Endusers should determine optimal dilutions

for their applications.



Detection of c-myc-tagged protein in 200, 100, and 50ng of E. coli lysate containing tagged fusion protein with #18826P diluted 1:25,000.

#### **DILUTION INSTRUCTIONS**

Dilute in PBS or medium that is identical to that used in the assay system.

## STORAGE AND STABILITY

This antibody is stable for at least one (1) year at 2-8°C.

For in vitro investigational use only. Not for use in therapeutic or diagnostic procedures.