

STAT5A (Phospho-Ser780) Polyclonal Antibody

ORDERING INFORMATION

Catalog No.: 43049

Format: 100ul at 1.0mg/ml in PBS (without Mg²⁺ and Ca²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Affinity-purified on phosphopeptide; non-phosphopeptide-reactive antibodies were removed by chromatography on non-phosphorylated peptide.

BACKGROUND

Signal transducers and activators of transcription (STATs) are transcription factors that play important roles in signaling pathways induced by many different cytokines and growth factors that regulate cell growth and differentiation, the immune response, antiviral activity, and homeostasis. Their activation and mechanisms of action are examples of direct signaling to the nucleus by extracellular polypeptide ligands without the involvement of second messengers, tyrosine phosphorylation of transcription factors, or transcription factors that make direct contact with membrane-bound receptors and DNA. Since the discovery of STAT 1 and 2, the family has been expanded to seven members.

SPECIFICATION SUMMARY

Antigen: Peptide sequence that includes phosphorylation site of serine 780 (R-L-S(p)-P-P) derived from human STAT5A and conjugated to KLH.


Accession no.: P42229, NP_003143.2

Host Species: Rabbit


Specificity: This antibody detects endogenous human, mouse, and rat STAT5A only when phosphorylated at serine 780.

APPLICATION

Immunoblotting: use at dilution of 1:500-1:1,000. **Immunohistochemistry:** use at dilution of 1:50-1:100.



Detection of STAT5A (phospho-Ser780) in extracts of K562 cells untreated or treated with IL-4.



Detection of STAT5A (phospho-Ser780) in paraffin-embedded human breast carcinoma tissue.

These are recommended working dilutions. Enduser should determine optimal dilutions for their applications.

STAT5A (Phospho-Ser780) Polyclonal Antibody

DILUTION INSTRUCTIONS

Dilute in PBS or medium that is identical to that used in the assay system.

STORAGE AND STABILITY

This antibody is stable for at least one (1) year at -20°C. Can be stored at 4°C for short-term use.

For in vitro investigational use only. Not intended for therapeutic or diagnostic applications.